

ODE/PDE Spring 2003 Qualifying Exam

Instruction: Complete all four problems. Clear and concise answers will improve your score.

1. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ denote a C^1 function and consider the system

$$\frac{du}{dt} = f(u)$$

for a vector function $u = u(t)$.

- a) What is a fixed point (in other terminology: critical point) of the system?
- b) When is a fixed point called stable in the sense of Lyapunov?
- c) Let u^* be a fixed point. How can one obtain stability or instability (in the sense of Lyapunov) using the eigenvalues of a certain matrix? (You do not have to prove the criteria.)
- d) Given the system

$$\begin{aligned}u'_1 &= u_1(1 - u_2) \\u'_2 &= u_2(4 - u_1)\end{aligned}$$

Determine all fixed points and discuss their stability.

2. a) Solve the scalar initial value problem

$$\begin{aligned}u'(t) &= |u(t)|, \quad t \geq 0, \\u(0) &= a,\end{aligned}$$

explicitly. Is the solution $u = u(t, a)$ everywhere differentiable as a function of $a \in \mathbb{R}$?

- b) Consider a scalar initial value problem

$$\begin{aligned}u'(t) &= f(u(t), t), \quad t \geq 0, \\u(0) &= a,\end{aligned}$$

where $f : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}$ is a given function. State conditions (without proof) which guarantee that the problem has a unique solution $u = u(t, a)$ existing for all time $t \geq 0$. Give conditions so that the partial derivative

$$u_a(t, a) = \frac{\partial u(t, a)}{\partial a}$$

exists and is continuous. Then prove the derivative is positive.

3. If $u = u(t) = u(x, t)$ then an initial boundary value problem (IBVP) for u is

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad (1)$$

$$u(x, 0) = f(x), \quad u(0, t) = 0, \quad u(1, t) = 0, \quad (2)$$

where $0 \leq x \leq 1$ and $t \geq 0$ and $f = f(x)$ is a given continuous function on $[0, 1]$.

- (a) Use separation of variables and Fourier series to give a formula for the solution u of the IBVP (1) and (2) in terms of the initial data f .
- (b) Is (1) elliptic, parabolic or hyperbolic? Why? If there is a maximum principle for this class of equations, state it. If this equation has characteristics, what are they? If $f(x) \equiv \sin(2\pi x)$, where is the maximum value of $u(x, t)$ for $0 \leq x \leq 1$ and $t \geq 0$?
- (c) Prove that the solution is infinitely differentiable for $t > 0$.
- (d) If $f(x) \equiv 1$, is the Fourier series solution continuous at $(x, t) = (0, 0)$?
- (e) If

$$\int_0^1 f(x) \sin(\pi x) dx = 0$$

what does the solution $u(x, t)$ look like for moderately large times? Give a formula and a sketch.

- (f) The solution of the IBVP can be written in the form

$$u(x, t) = \int_0^1 G(x, y, t) f(y) dy,$$

where G is the Green's function for the IBVP. Give a formula for G .

- (g) $F = F(x, t)$ is a continuous function on $t \geq 0, 0 \leq x \leq 1$, use Duhamel's principle to solve

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + F$$

along with the boundary conditions (2). Use the Green's function.

4. If $u = u(t) = u(x, t)$ then an initial value problem (IVP) for u is

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}, \quad (3)$$

$$u(x, 0) = f(x), \quad u_t(x, 0) = g(x) \quad (4)$$

where $t \geq 0$, $-\infty < x < \infty$, $u_t = \partial u / \partial t$ and $f = f(x)$ and $g = g(x)$ are a given continuous functions with compact support.

- (a) Derive d'Alembert's solution to (3) by first changing the equation to the variables $\xi = x + t$, $\eta = x - t$ and verifying that the resulting equation has solutions of the form $A(\xi) + B(\eta)$. Transform these solutions back to the x, t coordinates and then find a solution that satisfies the initial conditions.
- (b) Is (3) elliptic, parabolic or hyperbolic? Why? If there is a maximum principle for this class of equations, state it. If this equation has characteristics, what are they?
- (c) If

$$\chi_{[a,b]}(x) = \begin{cases} 1 & \text{if } a \leq x \leq b \\ 0 & \text{otherwise,} \end{cases}$$

is the characteristic function of the interval $[a, b]$, and if $f(x) = \chi_{[-1,1]}(x)$ and $g(x) \equiv 0$, then what is d'Alembert's solution to the IVP? Make a sketch in the (x, t) -plane of where this solution is non-zero. Is this solution continuous? Differentiable?

- (d) State reasonable conditions on f and g so that d'Alembert's solution to (3) and (4) is a *classical* solution. Is the solution with $f(x) = \chi_{[-1,1]}(x)$ and $g(x) \equiv 0$ a classical solution?
- (e) The energy $E = E(t)$ for the wave equation is the sum of the kinetic and potential energies:

$$E(t) = \frac{1}{2} \int_{-\infty}^{\infty} (u_t^2(x, t) + u_x^2(x, t)) \, dx.$$

Show that $E(t)$ is constant.

- (f) Show that the IVP (3) and (4) is well posed.
- (g) The solution of the IBVP can be written in the form

$$u(x, t) = \int_0^1 G(x, y, t) f(y) \, dy + \int_0^1 H(x, y, t) g(y) \, dy,$$

where G and H are Green's functions for the IBVP. Give formulas for G and H . How are G and H related?